

Quantum Mechanics  
Qualifying Exam - January 2017

*Notes and Instructions*

- There are 6 problems. Attempt them all as partial credit will be given.
- Write on only one side of the paper for your solutions.
- Write your alias on the top of every page of your solutions.
- Number each page of your solution with the problem number and page number (e.g. Problem 3, p. 2/4 is the second of four pages for the solution to problem 3.)
- You must show your work to receive full credit.

**Possibly useful formulas:**

Spin Operator

$$\vec{S} = \frac{\hbar}{2}\vec{\sigma}, \quad \sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (1)$$

In spherical coordinates,

$$\nabla^2\psi = \frac{1}{r} \frac{\partial^2}{\partial r^2} r\psi + \frac{1}{r^2 \sin\theta} \frac{\partial}{\partial\theta} (\sin\theta \frac{\partial\psi}{\partial\theta}) + \frac{1}{r^2 \sin^2\theta} \frac{\partial^2}{\partial\phi^2} \psi. \quad (2)$$

Harmonic oscillator wave functions

$$u_0(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-\frac{m\omega x^2}{2\hbar}}$$

$$u_1(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \sqrt{\frac{2m\omega}{\hbar}} x e^{-\frac{m\omega x^2}{2\hbar}}$$

Spherical Harmonics:

$$\begin{aligned} Y_{0,0}(\theta, \phi) &= \frac{1}{\sqrt{4\pi}} & Y_{2,0}(\theta, \phi) &= \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1) \\ Y_{1,0}(\theta, \phi) &= \sqrt{\frac{3}{4\pi}} \cos \theta & Y_{2,\pm 1}(\theta, \phi) &= \mp \sqrt{\frac{15}{8\pi}} e^{\pm i\phi} \cos \theta \sin \theta \\ Y_{1,\pm 1}(\theta, \phi) &= \mp \sqrt{\frac{3}{8\pi}} e^{\pm i\phi} \sin \theta & Y_{2,\pm 2}(\theta, \phi) &= \sqrt{\frac{15}{32\pi}} e^{\pm 2i\phi} \sin^2 \theta \end{aligned}$$

## Problem 1: Harmonic Oscillator (10 Points)

Consider the quantum mechanical simple harmonic oscillator.

- a. Using the raising and lower operators,  $\hat{a}$  and  $\hat{a}^\dagger$  find the average value of  $X$  and  $P$  for the state  $|n\rangle$ . **(1 Points)**
- b. Using the raising and lower operators,  $\hat{a}$  and  $\hat{a}^\dagger$  find the average value of  $X^2$  and  $P^2$  for the state  $|n\rangle$ . **(2 Points)**
- c. Using the raising and lower operators,  $\hat{a}$  and  $\hat{a}^\dagger$  find the root mean square deviations of  $X$  and  $P$  for the state  $|n\rangle$ . **(2 Points)**
- d. Find the uncertainty product for the state  $|n\rangle$  **(2 Points)**
- e. Find the average potential energy and average kinetic energy for the oscillator when it is in state  $|n\rangle$  **(3 Points)**

## Problem 2: Variational Method (10 Points)

The Hamiltonian of a one-dimensional harmonic oscillator is

$$H = \frac{P^2}{2m} + \frac{m\omega^2 X^2}{2}.$$

The ground state energy is  $E_0 = \hbar\omega/2$ .

Let us employ the variational method with the following trial function as the ground-state wave function

$$\langle x|\psi\rangle = \psi(x) = Ne^{-\beta|x|}.$$

- a. Determine the constant  $N$  by applying the normalization condition. (2 points)
- b. Find the value of  $\beta$  that minimizes  $\langle\psi|H|\psi\rangle$ . (2 points)
- c. What is the ground-state energy calculated with the variational method? (5 points)  
**N.B.** *The derivative of the trial function has a discontinuity.*
- d. How close do you get to the true ground-state energy? (1 points)

### Problem 3: Angular Momentum Hamiltonian (10 points)

Consider the following Hamiltonian for a spinless particle with orbital angular momentum  $\ell=2$ .

$$\hat{H} = \frac{3a}{2\hbar}\hat{L}_z - \frac{a}{\hbar^2}(\hat{L}_x^2 + \hat{L}_y^2)$$

where  $a$  is a constant greater than 0 and  $\hat{L}_i$  denotes the  $i^{\text{th}}$  component of the angular momentum operator.

- a) Calculate the energy spectrum of this Hamiltonian (2 pts)
- b) Suppose a particle with this Hamiltonian has the wavefunction

$$\Psi(\theta, \phi) = A(\sin \theta \cos \theta \cos \phi + \sin^2 \theta \sin \phi \cos \phi)$$

where  $\theta$  is the polar angle,  $\phi$  is the azimuthal angle, and  $A$  is a normalization constant. What is the average energy obtained in energy measurements on an ensemble of particles described by the wavefunction above? (3 pts)

- c) Assume the particle is in the lowest energy state (with  $\ell=2$ ) for  $t < 0$ . Starting at  $t=0$ , an external magnetic field is applied with

$$\hat{V}(t) = \frac{\lambda}{\hbar}\hat{L}_x e^{-t/\tau}$$

where  $\tau$  is the decay constant and  $\lambda$  is a constant. Calculate the transition probabilities to possible excited states after a very long time ( $\tau \ll t \rightarrow \infty$ ) using first order time-dependent perturbation theory. (5 pts)

## Problem 4: Hydrogen Atom (10 points)

Schrodinger's equation in spherical coordinates where the potential is only a function of  $r$  can be solved by using separation of variables:  $\Psi(r, \theta, \phi) = R(r)Y(\theta, \phi)$ .

a) In units  $2m = 1$  and  $\hbar = 1$ , show that using the change of variables  $u(r) \equiv rR(r)$ , one can obtain the radial Schrodinger's equation for the hydrogen atom. (1 pt)

$$\left[-\frac{d^2}{dr^2} - \frac{g^2}{r} + \frac{\ell(\ell+1)}{r^2}\right]u(r) = \epsilon u(r)$$

where  $g^2$  is the Coulomb strength and  $\epsilon$  is the energy.

b) The lowest eigenstate of a given  $\ell$  is known to have the form

$$u_\ell^0 = C_\ell r^{\ell+1} \exp(-r/a_\ell)$$

For a given  $\ell$ , determine the eigenvalue  $\epsilon_\ell^0$  and the size parameter  $a_\ell$ , in terms of  $g^2$  (2 pts).

Consider that the initial 3-dimensional wave function at time  $t=0$  is a superposition of the above states

$$\Psi(r, 0) = D(e^{-g^2 \frac{r}{2}} + g^2 r e^{-g^2 \frac{r}{4}} \cos \theta)$$

c) Determine  $\Psi(r, t)$  (1 pt)

d) Determine  $\langle \cos \theta \rangle$  as a function of time (3 pts).

e) Consider the hydrogen atom. Determine the most probable value of  $r$  for the ground state. (1 pt)

f) Consider a hydrogen atom placed in a weak constant uniform external electric field. Determine how the energy levels shift for the  $n=2$  state of hydrogen due to the electric field. (2 pts)

### Problem 5: 1/x potential (10 points)

An electron moves in one dimension and is confined to the right half space ( $x > 0$ ) where it has potential energy

$$V(x) = -\frac{e^2}{4x}$$

where  $e$  is the charge on an electron.

- a) What is the solution of Schrodinger's equation at large  $x$ ? (2 pts)
- b) What are the necessary boundary conditions (1 pt)
- c) Using the results of part a) and b), determine the ground state solution of the equation. (3 pts)
- d) Determine the ground state energy (2 pts)
- e) Find the expectation value  $\langle x \rangle$  in the ground state (2 pts)

## Problem 6: Measurements and Probability (10 points)

---

A three-level quantum system has a non-degenerate ground state and a two-fold degenerate excited state, defined by:

$$H|0\rangle = 0, \quad H|a\rangle = \epsilon|a\rangle, \quad H|b\rangle = \epsilon|b\rangle$$

where  $\epsilon$  is a positive constant energy.

- (a) (1 pt) Write down the matrix representation of  $H$  in the basis  $|0\rangle, |a\rangle, |b\rangle$ .
- (b) (2 pts.) Define the observable  $C$  by its operation on the eigenstates of  $H$ .

$$C|0\rangle = \gamma|a\rangle, \quad C|a\rangle = \gamma|0\rangle, \quad C|b\rangle = -\gamma|b\rangle \quad (3)$$

$\gamma > 0$ . What are all the possible outcomes of a measurement of  $C$ ?

- (c) (2 pts.) For each of the eigenstates of  $H$ , calculate the probability of measuring the different possible values for  $C$  if the system is in that eigenstate.
- (d) (1 pts.) Do  $H$  and  $C$  have common eigenstates? Are  $H$  and  $C$  compatible observables? Explain.
- (e) (2 pts.) At time  $t = 0$ , the system is in the eigenstate of  $C$  with the largest eigenvalue. Calculate the probabilities, as functions of time, of obtaining the different possible results of a measurement of  $C$ .
- (f) (2 pt.) At time  $t = 0$ , the system is in the state  $|\psi\rangle = \frac{1}{\sqrt{2}}(|a\rangle + |b\rangle)$ . Calculate the probabilities, as functions of time, of obtaining the different possible results of a measurement of  $C$ . Explain the differences in this result and what was found in part (e).